

# $K$ -theoretic Catalan functions

George H. Seelinger

*ghs9ae@virginia.edu*

University of Virginia

April 26, 2021

- Schubert calculus: connecting geometry and combinatorics
- Catalan functions: a new approach to old problems
- $K$ -theoretic Catalan functions

# Overview of Schubert Calculus Combinatorics

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of subvarieties in a variety  $X$ .

# Overview of Schubert Calculus Combinatorics

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of subvarieties in a variety  $X$ .



## Cohomology

Schubert basis  $\{\sigma_\lambda\}$  for  $H^*(X)$  with property  $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$

# Overview of Schubert Calculus Combinatorics

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of subvarieties in a variety  $X$ .



## Cohomology

Schubert basis  $\{\sigma_\lambda\}$  for  $H^*(X)$  with property  $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$



## Representatives

Special basis of polynomials  $\{f_\lambda\}$  such that  $f_\lambda \cdot f_\mu = \sum_\nu c_{\lambda\mu}^\nu f_\nu$

Combinatorial study of  $\{f_\lambda\}$  enlightens the geometry (and cohomology).

## Goal

Identify  $\{f_\lambda\}$  in explicit (simple) terms amenable to calculation and proofs.

# Algebra of Symmetric Functions

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ ?

# Algebra of Symmetric Functions

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ ?
- Symmetric polynomials ( $n = 3$ )

$$e_1 = x_1 + x_2 + x_3 \quad h_1 = x_1 + x_2 + x_3$$

$$e_2 = x_1x_2 + x_1x_3 + x_2x_3 \quad h_2 = x_1^2 + x_1x_2 + x_1x_3 + x_2^2 + x_2x_3 + x_3^2$$

$$e_3 = x_1x_2x_3 \quad h_3 = x_1^3 + x_1^2x_2 + x_1^2x_3 + x_1x_2^2 + \dots$$



# Algebra of Symmetric Functions

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ ?
- Symmetric polynomials ( $n = 3$ )

$$e_1 = x_1 + x_2 + x_3 \quad h_1 = x_1 + x_2 + x_3$$

$$e_2 = x_1x_2 + x_1x_3 + x_2x_3 \quad h_2 = x_1^2 + x_1x_2 + x_1x_3 + x_2^2 + x_2x_3 + x_3^2$$

$$e_3 = x_1x_2x_3 \quad h_3 = x_1^3 + x_1^2x_2 + x_1^2x_3 + x_1x_2^2 + \dots$$

- $\{f \in \mathbb{Q}[x_1, \dots, x_n] \mid \sigma.f = f \forall \sigma \in S_n\}$  forms a vector space,  $\Lambda_{\mathbb{Q}}$ .

# Algebra of Symmetric Functions

- Polynomials  $f \in \mathbb{Q}[x_1, \dots, x_n]$  satisfying  $\sigma.f = f$  for all  $\sigma \in S_n$ ?
- Symmetric polynomials ( $n = 3$ )

$$e_1 = x_1 + x_2 + x_3 \quad h_1 = x_1 + x_2 + x_3$$

$$e_2 = x_1x_2 + x_1x_3 + x_2x_3 \quad h_2 = x_1^2 + x_1x_2 + x_1x_3 + x_2^2 + x_2x_3 + x_3^2$$

$$e_3 = x_1x_2x_3 \quad h_3 = x_1^3 + x_1^2x_2 + x_1^2x_3 + x_1x_2^2 + \dots$$

- $\{f \in \mathbb{Q}[x_1, \dots, x_n] \mid \sigma.f = f \forall \sigma \in S_n\}$  forms a vector space,  $\Lambda_{\mathbb{Q}}$ .
- Bases indexed by integer partitions.

## Definition

$n \in \mathbb{Z}_{>0}$ , a *partition of  $n$*  is  $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell > 0)$  such that  $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = n$ .

## Definition

$n \in \mathbb{Z}_{>0}$ , a *partition* of  $n$  is  $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_\ell > 0)$  such that  $\lambda_1 + \lambda_2 + \dots + \lambda_\ell = n$ .

$$5 \rightarrow \begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}$$

$$4 + 1 \rightarrow \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}$$

$$3 + 2 \rightarrow \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}$$

$$3 + 1 + 1 \rightarrow \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}$$

$$2 + 2 + 1 \rightarrow \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \end{array}$$

$$2 + 1 + 1 + 1 \rightarrow \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}$$

$$1 + 1 + 1 + 1 + 1 \rightarrow \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \square \\ \hline \end{array}$$

## Definition

For  $m, n \in \mathbb{Z}_{>0}$ ,  $\text{Gr}(m, n) = \{V \subseteq \mathbb{C}^{m+n} \mid \dim V = m\}$ .

## Definition

For  $m, n \in \mathbb{Z}_{>0}$ ,  $\text{Gr}(m, n) = \{V \subseteq \mathbb{C}^{m+n} \mid \dim V = m\}$ .

- Topological structure of projective variety.

## Definition

For  $m, n \in \mathbb{Z}_{>0}$ ,  $\text{Gr}(m, n) = \{V \subseteq \mathbb{C}^{m+n} \mid \dim V = m\}$ .

- Topological structure of projective variety.
- $\text{Gr}(1, n) = \mathbb{C}P^n$

## Definition

For  $m, n \in \mathbb{Z}_{>0}$ ,  $\text{Gr}(m, n) = \{V \subseteq \mathbb{C}^{m+n} \mid \dim V = m\}$ .

- Topological structure of projective variety.
- $\text{Gr}(1, n) = \mathbb{C}P^n$
- “Schubert cell” decomposition

$$\text{Gr}(m, n) = \bigsqcup_{\lambda \subseteq (n^m)} \Omega_\lambda$$



## Definition

For  $m, n \in \mathbb{Z}_{>0}$ ,  $\text{Gr}(m, n) = \{V \subseteq \mathbb{C}^{m+n} \mid \dim V = m\}$ .

- Topological structure of projective variety.
- $\text{Gr}(1, n) = \mathbb{C}P^n$
- “Schubert cell” decomposition

$$\text{Gr}(m, n) = \bigsqcup_{\lambda \subseteq (n^m)} \Omega_\lambda$$

- Schubert varieties  $X_\lambda = \overline{\Omega_\lambda}$ .

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of Schubert varieties  $\{X_\lambda\}_{\lambda \subseteq (n^m)}$  in variety  $X = \text{Gr}(m, n)$ .

# Classical Schubert Calculus

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of Schubert varieties  $\{X_\lambda\}_{\lambda \subseteq (nm)}$  in variety  $X = \text{Gr}(m, n)$ .



## Cohomology

Schubert basis  $\{\sigma_\lambda\}_{\lambda \subseteq (nm)}$  for  $H^*(X)$  with property  $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$

# Classical Schubert Calculus

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of Schubert varieties  $\{X_\lambda\}_{\lambda \subseteq (n^m)}$  in variety  $X = \text{Gr}(m, n)$ .



## Cohomology

Schubert basis  $\{\sigma_\lambda\}_{\lambda \subseteq (n^m)}$  for  $H^*(X)$  with property  $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$



## Representatives

Special basis of Schur polynomials  $\{s_\lambda\}$  indexed by partitions such that  $s_\lambda \cdot s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$  for Littlewood-Richardson coefficients  $c_{\lambda\mu}^\nu$ .

## Example

*Semistandard tableaux*: columns increasing and rows non-decreasing.

$$T = \begin{array}{|c|c|c|c|} \hline 5 & & & \\ \hline 3 & 4 & & \\ \hline 2 & 3 & & \\ \hline 1 & 2 & 2 & 5 \\ \hline \end{array}$$
$$\begin{array}{|c|c|c|c|} \hline 8 & & & \\ \hline 7 & 9 & & \\ \hline 3 & 4 & & \\ \hline 1 & 2 & 5 & 6 \\ \hline \end{array}$$

## Example

*Semistandard tableaux*: columns increasing and rows non-decreasing.

$$T = \begin{array}{cccc} \boxed{5} & & & \\ \boxed{3} & \boxed{4} & & \\ \boxed{2} & \boxed{3} & & \\ \boxed{1} & \boxed{2} & \boxed{2} & \boxed{5} \end{array}$$
$$\begin{array}{cccc} \boxed{8} & & & \\ \boxed{7} & \boxed{9} & & \\ \boxed{3} & \boxed{4} & & \\ \boxed{1} & \boxed{2} & \boxed{5} & \boxed{6} \end{array}$$

$$\text{wt}(T) = (1, 3, 2, 1, 2) \quad (1, 1, 1, 1, 1, 1, 1, 1, 1)$$

## Example

*Semistandard tableaux*: columns increasing and rows non-decreasing.

$$T = \begin{array}{|c|c|c|c|} \hline 5 & & & \\ \hline 3 & 4 & & \\ \hline 2 & 3 & & \\ \hline 1 & 2 & 2 & 5 \\ \hline \end{array}$$

$$\text{wt}(T) = (1, 3, 2, 1, 2) \quad (1, 1, 1, 1, 1, 1, 1, 1, 1)$$

$$x^{\text{wt}(T)} = x_1^1 x_2^3 x_3^2 x_4^1 x_5^2 \quad x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8 x_9$$

$$\begin{array}{|c|c|c|c|} \hline 8 & & & \\ \hline 7 & 9 & & \\ \hline 3 & 4 & & \\ \hline 1 & 2 & 5 & 6 \\ \hline \end{array}$$

## Example

*Semistandard tableaux*: columns increasing and rows non-decreasing.

$$T = \begin{array}{|c|c|c|c|} \hline 5 & & & \\ \hline 3 & 4 & & \\ \hline 2 & 3 & & \\ \hline 1 & 2 & 2 & 5 \\ \hline \end{array} \qquad \begin{array}{|c|c|c|c|} \hline 8 & & & \\ \hline 7 & 9 & & \\ \hline 3 & 4 & & \\ \hline 1 & 2 & 5 & 6 \\ \hline \end{array}$$

$$\text{wt}(T) = (1, 3, 2, 1, 2) \quad (1, 1, 1, 1, 1, 1, 1, 1, 1)$$

$$x^{\text{wt}(T)} = x_1^1 x_2^3 x_3^2 x_4^1 x_5^2 \quad x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_8 x_9$$

$\text{SSYT}(\lambda) =$  all semistandard tableaux of shape  $\lambda$ .

$$\begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array}$$



# Schur functions $s_\lambda$

Schur function  $s_\lambda$  is a “weight generating function” of semistandard tableaux:

$$\begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array}$$

$$s_{\square}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

# Schur functions $s_\lambda$

Schur function  $s_\lambda$  is a “weight generating function” of semistandard tableaux:

$$\begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array}$$

$$s_{\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

$$s_\lambda(x) = \sum_{T \in \text{SSYT}(\lambda)} x^{\text{wt}(T)}$$

# Schur functions $s_\lambda$

Schur function  $s_\lambda$  is a “weight generating function” of semistandard tableaux:

$$\begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 1 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 1 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 2 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 2 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 2 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & \\ \hline 1 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & \\ \hline 1 & 2 \\ \hline \end{array}$$

$$s_{\square}(x_1, x_2, x_3) = x_1^2 x_2 + x_1^2 x_3 + x_2^2 x_3 + x_1 x_2^2 + x_1 x_3^2 + x_2 x_3^2 + 2x_1 x_2 x_3$$

$$s_\lambda(x) = \sum_{T \in \text{SSYT}(\lambda)} x^{\text{wt}(T)}$$

$s_\lambda(x)$  is homogeneous of degree  $\lambda_1 + \cdots + \lambda_\ell$ .

# Schur functions $s_\lambda$ (cont.)

## Pieri rule

Determines multiplicative structure:

$$s_r s_\lambda = \sum (1 \text{ or } 0) s_\nu$$

$$s_{\square} s_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} = s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}}$$

# Schur functions $s_\lambda$ (cont.)

## Pieri rule

Determines multiplicative structure:

$$s_r s_\lambda = \sum (1 \text{ or } 0) s_\nu$$

$$s_{\square} s_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} = s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}}$$

Iterate Pieri rule

$$s_{\mu_1} \cdots s_{\mu_r} s_\lambda = \sum (\# \text{ known tableaux}) s_\nu$$

## Pieri rule

Determines multiplicative structure:

$$s_r s_\lambda = \sum (1 \text{ or } 0) s_\nu$$

$$s_{\square} s_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} = s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}}$$

Iterate Pieri rule

$$s_{\mu_1} \cdots s_{\mu_r} s_\lambda = \sum (\# \text{ known tableaux}) s_\nu$$

Since  $s_{\mu_1} \cdots s_{\mu_r} = s_{(\mu_1, \dots, \mu_r)} + \text{lower order terms}$ , subtract to get

$$s_{(\mu_1, \dots, \mu_r)} s_\lambda = \sum c_{\lambda\mu}^\nu s_\nu$$

for well-understood *Littlewood-Richardson coefficients*  $c_{\lambda\mu}^\nu$ .

## Upshot

Let  $\{f_\lambda\}$  be a basis of  $\Lambda$  such that

- 1  $f_r = s_r$  and
- 2  $f_r f_\lambda$  satisfies the Pieri rule.

Then,  $f_\lambda = s_\lambda$ .

## Upshot

Let  $\{f_\lambda\}$  be a basis of  $\Lambda$  such that

- 1  $f_r = s_r$  and
- 2  $f_r f_\lambda$  satisfies the Pieri rule.

Then,  $f_\lambda = s_\lambda$ .

## Looking Ahead

This type of technique will be useful for establishing the equivalence of new formulas for other bases.



When examining Schubert representatives in  $\Lambda$ , we ask

When examining Schubert representatives in  $\Lambda$ , we ask

- Does it have a Pieri rule? ( $s_r s_\lambda = \sum s_\nu$ )

When examining Schubert representatives in  $\Lambda$ , we ask

- Does it have a Pieri rule? ( $s_r s_\lambda = \sum s_\nu$ )
- Does it have a direct formula? ( $s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} x^T$ )

There are many variations on classical Schubert calculus of the Grassmannian (Type A).

There are many variations on classical Schubert calculus of the Grassmannian (Type A).


Theory	$f_\lambda$
(Co)homology of Grassmannian	Schur functions
(Co)homology of flag variety	Schubert polynomials
Quantum cohomology of flag variety	Quantum Schuberts
(Co)homology of Types BCD Grassmannian	Schur- $P$ and $Q$ functions
(Co)homology of affine Grassmannian	(dual) $k$ -Schur functions
$K$ -theory of Grassmannian	Grothendieck polynomials
$K$ -homology of affine Grassmannian	$K$ - $k$ -Schur functions

There are many variations on classical Schubert calculus of the Grassmannian (Type A).

Theory	$f_\lambda$
(Co)homology of Grassmannian	Schur functions
(Co)homology of flag variety	Schubert polynomials
Quantum cohomology of flag variety	Quantum Schuberts
(Co)homology of Types BCD Grassmannian	Schur- $P$ and $Q$ functions
(Co)homology of affine Grassmannian	(dual) $k$ -Schur functions
$K$ -theory of Grassmannian	Grothendieck polynomials
$K$ -homology of affine Grassmannian	$K$ - $k$ -Schur functions

And many more!

# Big Picture



# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).



# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1^r s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$

# $k$ -Schur functions

- $s_\lambda^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_{1^r} s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$
- $s_\lambda^{(k)} = s_\lambda$  as  $k \rightarrow \infty$ .

# $k$ -Schur functions

- $s_\lambda^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1 r s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$
- $s_\lambda^{(k)} = s_\lambda$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}^{(2)} = s_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}^{(3)} + \underbrace{s_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}} + s_{\begin{smallmatrix} \square & \square & \square & \square \end{smallmatrix}}}_{s_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}^{(3)}}$$

# $k$ -Schur functions

- $s_\lambda^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1^r s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$
- $s_\lambda^{(k)} = s_\lambda$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{(2)}^{(2)} = s_{(2)}^{(3)} + s_{(3)}^{(3)} + s_{(4)}^{(3)}$$

- (Lam et al., 2010) gives geometric interpretation,

# $k$ -Schur functions

- $s_\lambda^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1^r s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$
- $s_\lambda^{(k)} = s_\lambda$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{(2)}^{(2)} = s_{(2)}^{(3)} + s_{(3)}^{(3)} + s_{(4)}^{(3)}$$

- (Lam et al., 2010) gives geometric interpretation,
- but no combinatorial interpretation of branching coefficients.

# $k$ -Schur functions

- $s_\lambda^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_{1^r} s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$
- $s_\lambda^{(k)} = s_\lambda$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{(2)}^{(2)} = s_{(2)}^{(3)} + s_{(1,1)}^{(3)} + s_{(1,1,1)}^{(3)}$$

The diagram illustrates the branching of the Schur function  $s_{(2)}^{(2)}$  into three Schur functions of degree 3. On the left is a 2x2 square Young diagram labeled  $s_{(2)}^{(2)}$ . This is equal to the sum of three Young diagrams: a 2x2 square labeled  $s_{(2)}^{(3)}$ , a Young diagram with two rows (1,1) labeled  $s_{(1,1)}^{(3)}$ , and a Young diagram with three rows (1,1,1) labeled  $s_{(1,1,1)}^{(3)}$ . Brackets are used to group the three diagrams on the right as a single sum.

- (Lam et al., 2010) gives geometric interpretation,
- but no combinatorial interpretation of branching coefficients.
- Branching with  $t$  important for Macdonald polynomial positivity.

# $k$ -Schur functions

- $s_\lambda^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1 r s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$
- $s_\lambda^{(k)} = s_\lambda$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{(2)}^{(2)} = s_{(2)}^{(3)} + s_{(3)}^{(3)} + s_{(4)}^{(3)}$$

The diagram illustrates the branching of the Schur function  $s_{(2)}^{(2)}$  into three Schur functions of degree 3. On the left is the Young diagram for  $s_{(2)}^{(2)}$ , a 2x2 square. This is equal to the sum of three Young diagrams:  $s_{(2)}^{(3)}$  (a 2x2 square),  $s_{(3)}^{(3)}$  (a 1x3 row), and  $s_{(4)}^{(3)}$  (a 1x4 row). Brackets are used to group the terms: a bracket under the first two terms is labeled  $s_{(2)}^{(3)}$ , and a bracket under the last two terms is labeled  $s_{(3)}^{(3)}$ .

- (Lam et al., 2010) gives geometric interpretation,
- but no combinatorial interpretation of branching coefficients.
- Branching with  $t$  important for Macdonald polynomial positivity.
- Many conjecturally equivalent definitions.



- Schubert calculus
- **Catalan functions: a new approach to old problems**
- *K*-theoretic Catalan functions

# Why a new definition of $k$ -Schur?

# Why a new definition of $k$ -Schur?

## Answer

- 1 (Blasiak et al., 2019) gives a new definition of  $s_\lambda^{(k)}$  and shows it is equivalent to many other previous definitions.

# Why a new definition of $k$ -Schur?

## Answer

- 1 (Blasiak et al., 2019) gives a new definition of  $s_\lambda^{(k)}$  and shows it is equivalent to many other previous definitions.
- 2 From a new definition, (Blasiak et al., 2019) shows the branching coefficients  $b_{\lambda\mu}$  in the expansion  $s_\lambda^{(k)} = \sum_\mu b_{\lambda\mu} s_\mu^{(k+1)}$  have combinatorial interpretation!

Key:

# Why a new definition of $k$ -Schur?

## Answer

- 1 (Blasiak et al., 2019) gives a new definition of  $s_\lambda^{(k)}$  and shows it is equivalent to many other previous definitions.
- 2 From a new definition, (Blasiak et al., 2019) shows the branching coefficients  $b_{\lambda\mu}$  in the expansion  $s_\lambda^{(k)} = \sum_\mu b_{\lambda\mu} s_\mu^{(k+1)}$  have combinatorial interpretation!

Key:  $\{s_\lambda^{(k)}\}_\lambda \subseteq \text{Catalan functions} = \text{large class of symmetric functions.}$

- Raising operators

# Ingredients for Catalan functions

- Raising operators
- Symmetric functions indexed by integer vectors

# Ingredients for Catalan functions

- Raising operators
- Symmetric functions indexed by integer vectors
- Root ideals



# Raising Operators on Symmetric Functions

- Raising operators  $R_{i,j}$  act on diagrams

$$R_{1,3} \left( \begin{array}{|c|c|c|c|} \hline \color{red}{\square} & & & \\ \hline \square & & & \\ \hline \square & \square & \square & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline \square & & & \\ \hline \square & \square & \square & \color{red}{\square} \\ \hline \square & \square & \square & \square \\ \hline \end{array} \quad R_{2,3} \left( \begin{array}{|c|c|} \hline \square & \color{red}{\square} \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|c|} \hline \square & & \\ \hline \square & \square & \color{red}{\square} \\ \hline \square & \square & \\ \hline \end{array}$$

# Raising Operators on Symmetric Functions

- Raising operators  $R_{i,j}$  act on diagrams

$$R_{1,3} \left( \begin{array}{|c|c|c|c|} \hline \color{red}{\square} & & & \\ \hline \square & & & \\ \hline \square & \square & \square & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline \square & & & \\ \hline \square & \square & \square & \color{red}{\square} \\ \hline \square & \square & \square & \square \\ \hline \end{array} \quad R_{2,3} \left( \begin{array}{|c|c|} \hline \square & \color{red}{\square} \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|c|} \hline \square & & \\ \hline \square & \square & \color{red}{\square} \\ \hline \square & \square & \\ \hline \end{array}$$

- Extend action to a symmetric function  $f_\lambda$  by  $R_{i,j}(f_\lambda) = f_{\lambda+\epsilon_i-\epsilon_j}$ .

# Raising Operators on Symmetric Functions

- Raising operators  $R_{i,j}$  act on diagrams

$$R_{1,3} \left( \begin{array}{|c|c|c|c|} \hline \color{red}{\square} & & & \\ \hline \square & & & \\ \hline \square & \square & \square & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & \square & & \\ \hline \square & & & \\ \hline \square & \square & \square & \color{red}{\square} \\ \hline \end{array} \quad R_{2,3} \left( \begin{array}{|c|c|} \hline \square & \color{red}{\square} \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|c|} \hline & \square & \\ \hline \square & & \color{red}{\square} \\ \hline \square & \square & \\ \hline \end{array}$$

- Extend action to a symmetric function  $f_\lambda$  by  $R_{i,j}(f_\lambda) = f_{\lambda+\epsilon_i-\epsilon_j}$ .
- For  $h_\lambda = s_{\lambda_1} \cdots s_{\lambda_r}$ , we have the *Jacobi-Trudi identity*

$$s_\lambda = \prod_{i < j} (1 - R_{ij}) h_\lambda$$

# Raising Operators on Symmetric Functions

- Raising operators  $R_{i,j}$  act on diagrams

$$R_{1,3} \left( \begin{array}{|c|c|c|c|} \hline \color{red}{\square} & & & \\ \hline \square & & & \\ \hline \square & \square & \square & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & \square & & \\ \hline \square & & & \\ \hline \square & \square & \square & \color{red}{\square} \\ \hline \end{array} \quad R_{2,3} \left( \begin{array}{|c|c|c|} \hline & & \color{red}{\square} \\ \hline \square & \square & \\ \hline \square & \square & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|} \hline & \square & \\ \hline \square & \square & \color{red}{\square} \\ \hline \square & \square & \\ \hline \end{array}$$

- Extend action to a symmetric function  $f_\lambda$  by  $R_{i,j}(f_\lambda) = f_{\lambda+\epsilon_i-\epsilon_j}$ .
- For  $h_\lambda = s_{\lambda_1} \cdots s_{\lambda_r}$ , we have the *Jacobi-Trudi identity*

$$s_\lambda = \prod_{i < j} (1 - R_{ij}) h_\lambda$$

$$s_{22} = (1 - R_{12}) h_{22} = h_{22} - h_{31}$$

$$s_{211} = (1 - R_{12})(1 - R_{23})(1 - R_{13}) h_{211}$$

$$= h_{211} - h_{301} - h_{220} - \color{red}{h_{310}} + \color{red}{h_{310}} + \underbrace{h_{32-1}}_{=0} + h_{400} - \underbrace{h_{41-1}}_{=0}$$

some terms cancel

# Raising Operators on Symmetric Functions

Upside: gives definition for Schur function indexed by any integer vector  $\alpha \in \mathbb{Z}^\ell$ .

# Raising Operators on Symmetric Functions

Upside: gives definition for Schur function indexed by any integer vector  $\alpha \in \mathbb{Z}^\ell$ . Straightening:

$$s_\alpha = \prod_{i < j} (1 - R_{ij}) h_\alpha = \begin{cases} \pm s_\lambda & \text{for a partition } \lambda \\ 0 & \end{cases}$$

# Raising Operators on Symmetric Functions

Upside: gives definition for Schur function indexed by any integer vector  $\alpha \in \mathbb{Z}^\ell$ . Straightening:

$$s_\alpha = \prod_{i < j} (1 - R_{ij}) h_\alpha = \begin{cases} \pm s_\lambda & \text{for a partition } \lambda \\ 0 & \end{cases}$$

Simplifies formulas. E.g., for  $\langle s_{1^r}^\perp s_\lambda, s_\mu \rangle = \langle s_\lambda, s_{1^r} s_\mu \rangle$  (note  $\langle s_\lambda, s_\mu \rangle = \delta_{\lambda\mu}$ ),

$$s_{1^r}^\perp s_\lambda =$$

# Raising Operators on Symmetric Functions

Upside: gives definition for Schur function indexed by any integer vector  $\alpha \in \mathbb{Z}^\ell$ . Straightening:

$$s_\alpha = \prod_{i < j} (1 - R_{ij}) h_\alpha = \begin{cases} \pm s_\lambda & \text{for a partition } \lambda \\ 0 & \end{cases}$$

Simplifies formulas. E.g., for  $\langle s_{1^r}^\perp s_\lambda, s_\mu \rangle = \langle s_\lambda, s_{1^r} s_\mu \rangle$  (note  $\langle s_\lambda, s_\mu \rangle = \delta_{\lambda\mu}$ ),

$$s_{1^r}^\perp s_\lambda = \sum_{S \subseteq [1, \ell], |S|=r} s_{\lambda - \epsilon_S}$$

$$s_{1^2}^\perp s_{333} = s_{322} + s_{232} + s_{223}$$



# Raising Operators on Symmetric Functions

Upside: gives definition for Schur function indexed by any integer vector  $\alpha \in \mathbb{Z}^\ell$ . Straightening:

$$s_\alpha = \prod_{i < j} (1 - R_{ij}) h_\alpha = \begin{cases} \pm s_\lambda & \text{for a partition } \lambda \\ 0 & \end{cases}$$

Simplifies formulas. E.g., for  $\langle s_{1^r}^\perp s_\lambda, s_\mu \rangle = \langle s_\lambda, s_{1^r} s_\mu \rangle$  (note  $\langle s_\lambda, s_\mu \rangle = \delta_{\lambda\mu}$ ),


$$s_{1^r}^\perp s_\lambda = \sum_{S \subseteq [1, \ell], |S|=r} s_{\lambda - \epsilon_S}$$

$$s_{1^2}^\perp s_{333} = s_{322} + s_{232} + s_{223}$$

$$s_{1^3}^\perp s_{333} = s_{222}$$

# Root Ideals


A root ideal  $\Psi$  of type  $A_{\ell-1}$  positive roots: given by Dyck path (lattice path above diagonal).



$\Psi =$  Roots above Dyck path  
 $\Delta_{\ell}^{+} \setminus \Psi =$  Non-roots below

# Root Ideals

A root ideal  $\Psi$  of type  $A_{\ell-1}$  positive roots: given by Dyck path (lattice path above diagonal).



$\Psi =$  Roots above Dyck path  
 $\Delta_{\ell}^{+} \setminus \Psi =$  Non-roots below


Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)

For  $\Psi$  and  $\gamma \in \mathbb{Z}^{\ell}$

$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^{+} \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

# Root Ideals

A root ideal  $\Psi$  of type  $A_{\ell-1}$  positive roots: given by Dyck path (lattice path above diagonal).



$\Psi =$  Roots above Dyck path  
 $\Delta_{\ell}^{+} \setminus \Psi =$  Non-roots below

Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)


For  $\Psi$  and  $\gamma \in \mathbb{Z}^{\ell}$

$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^{+} \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

- $\Psi = \emptyset \implies H(\emptyset; \gamma) = s_{\gamma}$

# Root Ideals

A root ideal  $\Psi$  of type  $A_{\ell-1}$  positive roots: given by Dyck path (lattice path above diagonal).



$\Psi$  = Roots above Dyck path  
 $\Delta_{\ell}^{+} \setminus \Psi$  = Non-roots below

## Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)

For  $\Psi$  and  $\gamma \in \mathbb{Z}^{\ell}$

$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^{+} \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

- $\Psi = \emptyset \implies H(\emptyset; \gamma) = s_{\gamma}$
- $\Psi = \text{all roots} \implies H(\Psi; \gamma) = h_{\gamma}$

## Intuition

Catalan functions interpolate between  $h_\lambda$  and  $s_\lambda$ .

## Intuition

Catalan functions interpolate between  $h_\lambda$  and  $s_\lambda$ .

## Theorem (Blasiak et al., 2020)

For  $\Psi$  any root ideal and  $\lambda$  a partition,  $H(\Psi; \lambda)$  is Schur positive!  
Precisely,  $H(\Psi; \lambda) = \sum_\nu c'_{\Psi, \lambda} s_\nu$  satisfies  $c'_{\Psi, \lambda} \in \mathbb{Z}_{\geq 0}$ .

## $k$ -Schur root ideal for $\lambda$

$$\begin{aligned}\psi &= \Delta^k(\lambda) = \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$



# Catalan functions

## $k$ -Schur root ideal for $\lambda$

$$\begin{aligned}\psi = \Delta^k(\lambda) &= \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3					
	3				
		2			
			2		
				1	
					1

← row  $i$  has  $4 - \lambda_i$  non-roots

# Catalan functions

## $k$ -Schur root ideal for $\lambda$

$$\begin{aligned}\psi = \Delta^k(\lambda) &= \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3						
	3					
		2				
			2			
				1		
					1	

$\leftarrow$  row  $i$  has  $4 - \lambda_i$  non-roots

$k$ -Schur is a Catalan function (Blasiak et al., 2019).

For partition  $\lambda$  with  $\lambda_1 \leq k$ ,

$$s_\lambda^{(k)} = H(\Delta^k(\lambda); \lambda).$$

## Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof:  $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof:  $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3					
	3				
		2			
			2		
				1	
					1

$$\Delta^5(4, 4, 3, 3, 2, 2) =$$

4					
	4				
		3			
			3		
				2	
					2

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof:  $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3					
	3				
		2			
			2		
				1	
					1

$$\Delta^5(4, 4, 3, 3, 2, 2) =$$

4					
	4				
		3			
			3		
				2	
					2

Pieri:

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = \sum_\mu a_{\lambda+1^\ell, \mu} s_\mu^{(k+1)}$$

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof:  $k - \lambda_i = (k + 1) - (\lambda_i + 1)$


$$\Delta^4(3, 3, 2, 2, 1, 1) = \begin{array}{|c|c|c|c|c|c|} \hline 3 & & & & & \\ \hline & 3 & & & & \\ \hline & & 2 & & & \\ \hline & & & 2 & & \\ \hline & & & & 1 & \\ \hline & & & & & 1 \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array} \quad \Delta^5(4, 4, 3, 3, 2, 2) = \begin{array}{|c|c|c|c|c|c|} \hline 4 & & & & & \\ \hline & 4 & & & & \\ \hline & & 3 & & & \\ \hline & & & 3 & & \\ \hline & & & & 2 & \\ \hline & & & & & 2 \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}$$

Branching is a special case of Pieri:

$$s_\lambda^{(k)} = s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = \sum_\mu a_{\lambda+1^\ell, \mu} s_\mu^{(k+1)}$$



# Big Picture



- Schubert calculus
- Catalan functions: a new approach to old problems
- ***K*-theoretic Catalan functions**

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda + \text{lower degree terms}$ .

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda +$  lower degree terms.
- Satisfies Pieri rule on “set-valued strips”

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda +$  lower degree terms.
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} + g_{3211} - g_{42} - g_{33} - 2g_{321} + g_{32}$$



Add (addable) or mark (removable) in any combination of  $r$  boxes, but only once per row.

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda +$  lower degree terms.
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} + g_{3211} - g_{42} - g_{33} - 2g_{321} + g_{32}$$



Add (addable) or mark (removable) in any combination of  $r$  boxes, but only once per row.

- $g_\lambda = \prod_{i < j} (1 - R_{ij}) k_\lambda$  for  $k_\lambda$  and inhomogeneous analogue of  $h_\lambda$ .

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda +$  lower degree terms.
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} + g_{3211} - g_{42} - g_{33} - 2g_{321} + g_{32}$$



Add (addable) or mark (removable) in any combination of  $r$  boxes, but only once per row.

- $g_\lambda = \prod_{i < j} (1 - R_{ij}) k_\lambda$  for  $k_\lambda$  and inhomogeneous analogue of  $h_\lambda$ .
- Dual to Grothendieck polynomials  $G_\lambda$ : Schubert representatives for  $K^*(Gr(m, n))$

- Inhomogeneous basis:  $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$



- Inhomogeneous basis:  $g_\lambda^{(k)} = s_\lambda^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”
- A  $(k + 1)$ -core is a partition with no cell of hook length  $k + 1$ .



- Inhomogeneous basis:  $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”
- A  $(k + 1)$ -core is a partition with no cell of hook length  $k + 1$ .



$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions  $\leftrightarrow$  3-cores


# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_\lambda^{(k)} = s_\lambda^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”
- A  $(k + 1)$ -core is a partition with no cell of hook length  $k + 1$ .



$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions  $\leftrightarrow$  3-cores



# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_\lambda^{(k)} = s_\lambda^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”
- A  $(k + 1)$ -core is a partition with no cell of hook length  $k + 1$ .




$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions  $\leftrightarrow$  3-cores

# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_\lambda^{(k)} = s_\lambda^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”
- A  $(k + 1)$ -core is a partition with no cell of hook length  $k + 1$ .



Conjecture (Lam et al., 2010; Morse, 2011)

$g_{\lambda}^{(k)}$  have positive branching into  $g_{\mu}^{(k+1)}$ .

Conjecture (Lam et al., 2010; Morse, 2011)

$g_{\lambda}^{(k)}$  have positive branching into  $g_{\mu}^{(k+1)}$ .

Problem

No direct formula for  $g_{\lambda}^{(k)}$



## Solution

Find a formula for  $g_{\lambda}^{(k)}$  analogous to raising operator formula for  $s_{\lambda}^{(k)}$ .

## Solution

Find a formula for  $g_{\lambda}^{(k)}$  analogous to raising operator formula for  $s_{\lambda}^{(k)}$ .

Requires an inhomogeneous refinement of Catalan functions.

# An Extra Ingredient: Lowering Operators

Lowering Operators  $L_j(f_\lambda) = f_{\lambda - \epsilon_j}$

$$L_3 \left( \begin{array}{|c|c|c|c|} \hline \color{red}{\square} & & & \\ \hline \square & & & \\ \hline \square & \square & \square & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|c|} \hline \square & & \\ \hline \square & \square & \square \\ \hline \end{array}, \quad L_1 \left( \begin{array}{|c|c|c|c|} \hline \square & & & \\ \hline \square & & & \\ \hline \square & \square & \color{red}{\square} & \square \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array}$$

## $K$ -theoretic Catalan function

Let  $\Psi, \mathcal{L} \subseteq \Delta_\ell^+$  be order ideals of positive roots and  $\gamma \in \mathbb{Z}^\ell$ , then

$$K(\Psi; \mathcal{L}; \gamma) := \prod_{(i,j) \in \mathcal{L}} (1 - L_j) \prod_{(i,j) \in \Delta_\ell^+ \setminus \Psi} (1 - R_{ij}) k_\gamma$$

## $K$ -theoretic Catalan function

Let  $\Psi, \mathcal{L} \subseteq \Delta_\ell^+$  be order ideals of positive roots and  $\gamma \in \mathbb{Z}^\ell$ , then

$$K(\Psi; \mathcal{L}; \gamma) := \prod_{(i,j) \in \mathcal{L}} (1 - L_j) \prod_{(i,j) \in \Delta_\ell^+ \setminus \Psi} (1 - R_{ij}) k_\gamma$$

“ $\Psi$  =raising ideal,  $\mathcal{L}$  =lowering ideal.”

# Affine $K$ -Theory Representatives with Raising Operators

## $K$ -theoretic Catalan function

Let  $\Psi, \mathcal{L} \subseteq \Delta_\ell^+$  be order ideals of positive roots and  $\gamma \in \mathbb{Z}^\ell$ , then

$$K(\Psi; \mathcal{L}; \gamma) := \prod_{(i,j) \in \mathcal{L}} (1 - L_j) \prod_{(i,j) \in \Delta_\ell^+ \setminus \Psi} (1 - R_{ij}) k_\gamma$$

“ $\Psi$  =raising ideal,  $\mathcal{L}$  =lowering ideal.”

## Example

non-roots of  $\Psi$ , roots of  $\mathcal{L}$

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

$$\begin{aligned} K(\Psi; \mathcal{L}; 54332) \\ = (1 - L_4)^2 (1 - L_5)^2 (1 - R_{12}) (1 - R_{34}) (1 - R_{45}) k_{54332} \end{aligned}$$

Answer (Blasiak-Morse-S., 2020)

Answer (Blasiak-Morse-S., 2020)

For  $K$ -homology of affine Grassmannian,  $g_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$  since this family satisfies the Pieri rule.




# Affine $K$ -Theory Representatives with Raising Operators

Answer (Blasiak-Morse-S., 2020)

For  $K$ -homology of affine Grassmannian,  $g_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$  since this family satisfies the Pieri rule.

Example



$$\Delta_9^+ / \Delta^4(332111111), \Delta^5(332111111)$$

# Pieri Rule Illustrated (Recurrences)

A “graphical calculus.”

$$g_1 g_{211}^{(2)}$$

# Pieri Rule Illustrated (Recurrences)

A “graphical calculus.”

$$g_1 g_{211}^{(2)}$$

$$=$$

2						
	1					
		1				
			0			
				0		
					0	
						1

# Pieri Rule Illustrated (Recurrences)

A “graphical calculus.”

$$g_1 g_{211}^{(2)}$$

$$=$$

2						
	1					
		1				
			0			
				0		
					0	
						1

$$=$$

2						
	1					
		1				
			0			
				0		
					0	
						1




$$+$$

2						
	1					
		1				
			0			
				0		
						1

# Pieri Rule Illustrated (Recurrences)

A “graphical calculus.”

$$g_1 g_{211}^{(2)}$$



# Pieri Rule Illustrated (Straightening)

$$g_1 g_{211}^{(2)} =$$

2						
	1					
		1				
			0			
				0		
					0	
						1

$$+$$

2						
	1					
		1				
			0			
				0		
					0	
						1

$$+$$

2						
	1					
		1				
			0			
				0		
					0	
						1

# Pieri Rule Illustrated (Straightening)

$$g_1 g_{211}^{(2)} =$$

2						
	1					
		1				
			0			
				0		
					0	
						1

$$+$$

2						
	1					
		1				
			0			
				0		
					0	
						1

$$+$$

2						
	1					
		1				
			0			
				0		
					0	
						1

$$=$$

2			
	1		
		1	
			1

$$-$$

2		
	1	
		1

$$-$$

2		
	1	
		1

# Pieri Rule Illustrated (Straightening)


$$\begin{aligned}
 g_1 g_{211}^{(2)} &= \begin{array}{|c|c|c|c|c|c|} \hline 2 & & & & & \\ \hline & 1 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline & & & & & 1 \\ \hline \end{array} + \begin{array}{|c|c|c|c|c|c|} \hline 2 & & & & & \\ \hline & 1 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline & & & & & 1 \\ \hline \end{array} + \begin{array}{|c|c|c|c|c|c|} \hline 2 & & & & & \\ \hline & 1 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline & & & & & 1 \\ \hline \end{array} \\
 &= \begin{array}{|c|c|c|c|} \hline 2 & & & \\ \hline & 1 & & \\ \hline & & 1 & \\ \hline & & & 1 \\ \hline \end{array} - \begin{array}{|c|c|c|} \hline 2 & & \\ \hline & 1 & \\ \hline & & 1 \\ \hline \end{array} - \begin{array}{|c|c|c|} \hline 2 & & \\ \hline & 1 & \\ \hline & & 1 \\ \hline \end{array} \\
 &= g_{2111}^{(2)} - g_{211}^{(2)} - g_{211}^{(2)}
 \end{aligned}$$



# Pieri Rule Illustrated (Straightening)

$$\begin{aligned}
 g_1 g_{211}^{(2)} &= \begin{array}{|c|c|c|c|c|c|} \hline 2 & & & & & \\ \hline & 1 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline & & & & & 1 \\ \hline \end{array} + \begin{array}{|c|c|c|c|c|c|} \hline 2 & & & & & \\ \hline & 1 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline & & & & & 1 \\ \hline \end{array} + \begin{array}{|c|c|c|c|c|c|} \hline 2 & & & & & \\ \hline & 1 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 1 \\ \hline & & & & & \\ \hline \end{array} \\
 &= \begin{array}{|c|c|c|c|} \hline 2 & & & \\ \hline & 1 & & \\ \hline & & 1 & \\ \hline & & & 1 \\ \hline \end{array} - \begin{array}{|c|c|c|} \hline 2 & & \\ \hline & 1 & \\ \hline & & 1 \\ \hline \end{array} - \begin{array}{|c|c|c|} \hline 2 & & \\ \hline & 1 & \\ \hline & & 1 \\ \hline \end{array} \\
 &= g_{2111}^{(2)} - g_{211}^{(2)} - g_{211}^{(2)}
 \end{aligned}$$

3-core perspective:



Theorem (Blasiak-Morse-S., 2020)

## Theorem (Blasiak-Morse-S., 2020)

The  $g_\lambda^{(k)}$  are “shift invariant”, i.e. for  $\ell = \ell(\lambda)$

$$G_{1^\ell}^\perp g_{\lambda+1^\ell}^{(k+1)} = g_\lambda^{(k)}$$

## Theorem (Blasiak-Morse-S., 2020)

The  $g_\lambda^{(k)}$  are “shift invariant”, i.e. for  $\ell = \ell(\lambda)$

$$G_{1^\ell}^\perp g_{\lambda+1^\ell}^{(k+1)} = g_\lambda^{(k)}$$


## Theorem (Blasiak-Morse-S., 2020)

The branching coefficients in

$$g_\lambda^{(k)} = \sum_{\mu} a_{\lambda\mu} g_\mu^{(k+1)}$$

satisfy  $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$ .

# Big Picture



# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

Conjecture (Ikeda et al., 2018)

For  $w \in S_{k+1}$  and  $\mathfrak{G}_w^Q$  a “quantum Grothendieck polynomial”,

$$\Phi(\mathfrak{G}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in Des(w)} \tau_i}$$

# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

Conjecture (Ikeda et al., 2018)

For  $w \in S_{k+1}$  and  $\mathfrak{G}_w^Q$  a “quantum Grothendieck polynomial”,

$$\Phi(\mathfrak{G}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in Des(w)} \tau_i}$$

satisfies  $\tilde{g}_w = g_\lambda^{(k)} + \sum_{\mu} a_{\lambda\mu} g_\mu^{(k)}$  such that  $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$ .



# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

## Conjecture (Ikeda et al., 2018)

For  $w \in S_{k+1}$  and  $\mathfrak{G}_w^Q$  a “quantum Grothendieck polynomial”,

$$\Phi(\mathfrak{G}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in Des(w)} \tau_i}$$

satisfies  $\tilde{g}_w = g_\lambda^{(k)} + \sum_\mu a_{\lambda\mu} g_\mu^{(k)}$  such that  $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$ .

## Theorem (Blasiak-Morse-S., 2020)

If  $\lambda \subseteq (d^{k+1-d})$  for some  $1 \leq d \leq k$ , then  $g_\lambda^{(k)} = g_\lambda$ . Thus, conjecture is true for  $w$  a Grassmannian permutation (i.e.  $w$  has only one descent).

# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

## Conjecture (Ikeda et al., 2018)

For  $w \in S_{k+1}$  and  $\mathfrak{G}_w^Q$  a “quantum Grothendieck polynomial”,

$$\Phi(\mathfrak{G}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in Des(w)} \tau_i}$$

satisfies  $\tilde{g}_w = g_\lambda^{(k)} + \sum_{\mu} a_{\lambda\mu} g_\mu^{(k)}$  such that  $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$ .

## Theorem (Blasiak-Morse-S., 2020)

If  $\lambda \subseteq (d^{k+1-d})$  for some  $1 \leq d \leq k$ , then  $g_\lambda^{(k)} = g_\lambda$ . Thus, conjecture is true for  $w$  a Grassmannian permutation (i.e.  $w$  has only one descent).

## Conjecture (Blasiak-Morse-S., 2020)

$$\tilde{g}_w = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$$

Definition (Blasiak-Morse-S., 2020)

For any partition  $\lambda$  with  $\lambda_1 \leq k$ , we set

$$\tilde{g}_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$$

# Closed $K$ - $k$ -Schur functions

## Definition (Blasiak-Morse-S., 2020)

For any partition  $\lambda$  with  $\lambda_1 \leq k$ , we set

$$\tilde{g}_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$$

## Conjecture (Blasiak-Morse-S., 2020)

These  $\tilde{g}_\mu^{(k)}$  satisfy the following properties.

## Definition (Blasiak-Morse-S., 2020)

For any partition  $\lambda$  with  $\lambda_1 \leq k$ , we set

$$\tilde{g}_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$$

## Conjecture (Blasiak-Morse-S., 2020)

These  $\tilde{g}_\mu^{(k)}$  satisfy the following properties.

- The coefficients in  $G_{1^m}^\perp \tilde{g}_\mu^{(k)} = \sum_\nu c_{\mu\nu} \tilde{g}_\nu^{(k)}$  satisfy  $(-1)^{|\mu|-|\nu|} c_{\mu\nu} \in \mathbb{Z}_{\geq 0}$ .

## Definition (Blasiak-Morse-S., 2020)

For any partition  $\lambda$  with  $\lambda_1 \leq k$ , we set

$$\tilde{g}_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$$

## Conjecture (Blasiak-Morse-S., 2020)

These  $\tilde{g}_\mu^{(k)}$  satisfy the following properties.

- The coefficients in  $G_{1^m}^\perp \tilde{g}_\mu^{(k)} = \sum_\nu c_{\mu\nu} \tilde{g}_\nu^{(k)}$  satisfy  $(-1)^{|\mu|-|\nu|} c_{\mu\nu} \in \mathbb{Z}_{\geq 0}$ .
- The coefficients in  $\tilde{g}_\mu^{(k)} = \sum_\nu a_{\mu\nu} \tilde{g}_\nu^{(k+1)}$  satisfy  $(-1)^{|\mu|-|\nu|} a_{\mu\nu} \in \mathbb{Z}_{\geq 0}$ .

## Definition (Blasiak-Morse-S., 2020)

For any partition  $\lambda$  with  $\lambda_1 \leq k$ , we set

$$\tilde{g}_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$$

## Conjecture (Blasiak-Morse-S., 2020)

These  $\tilde{g}_\mu^{(k)}$  satisfy the following properties.

- The coefficients in  $G_{1^m}^\perp \tilde{g}_\mu^{(k)} = \sum_\nu c_{\mu\nu} \tilde{g}_\nu^{(k)}$  satisfy  $(-1)^{|\mu|-|\nu|} c_{\mu\nu} \in \mathbb{Z}_{\geq 0}$ .
- The coefficients in  $\tilde{g}_\mu^{(k)} = \sum_\nu a_{\mu\nu} \tilde{g}_\nu^{(k+1)}$  satisfy  $(-1)^{|\mu|-|\nu|} a_{\mu\nu} \in \mathbb{Z}_{\geq 0}$ .
- The coefficients in  $\tilde{g}_\mu^{(k)} = \sum_\nu b_{\mu\nu} g_\nu^{(k)}$  satisfy  $(-1)^{|\mu|-|\nu|} b_{\mu\nu} \in \mathbb{Z}_{\geq 0}$ .

## Theorem (S., 2021)

*For  $1 \leq d \leq k$ , set  $R_d = ((k + 1 - d)^d)$  to be the  $k$ -rectangle partition.*



## Theorem (S., 2021)

For  $1 \leq d \leq k$ , set  $R_d = ((k + 1 - d)^d)$  to be the  $k$ -rectangle partition. Then,

$$\tilde{g}_{R_d}^{(k)} \tilde{g}_\mu^{(k)} = \tilde{g}_{\mu \cup R_d}^{(k)},$$

where  $\mu \cup R_d$  is the partition given by sorting  $(\mu, R_d)$ .

## Theorem (S., 2021)

For  $1 \leq d \leq k$ , set  $R_d = ((k + 1 - d)^d)$  to be the  $k$ -rectangle partition. Then,

$$\tilde{g}_{R_d}^{(k)} \tilde{g}_{\mu}^{(k)} = \tilde{g}_{\mu \cup R_d}^{(k)},$$

where  $\mu \cup R_d$  is the partition given by sorting  $(\mu, R_d)$ .

- Must be true for geometric connection with Peterson isomorphism.

## Theorem (S., 2021)

For  $1 \leq d \leq k$ , set  $R_d = ((k + 1 - d)^d)$  to be the  $k$ -rectangle partition. Then,

$$\tilde{g}_{R_d}^{(k)} \tilde{g}_\mu^{(k)} = \tilde{g}_{\mu \cup R_d}^{(k)},$$

where  $\mu \cup R_d$  is the partition given by sorting  $(\mu, R_d)$ .

- Must be true for geometric connection with Peterson isomorphism.
- Corresponding result for  $s_\lambda^{(k)}$  is known, but this gives a Catalan/ $K$ -theoretic Catalan proof.

# $k$ -Rectangle Property

## Theorem (S., 2021)

For  $1 \leq d \leq k$ , set  $R_d = ((k + 1 - d)^d)$  to be the  $k$ -rectangle partition. Then,

$$\tilde{g}_{R_d}^{(k)} \tilde{g}_\mu^{(k)} = \tilde{g}_{\mu \cup R_d}^{(k)},$$

where  $\mu \cup R_d$  is the partition given by sorting  $(\mu, R_d)$ .

- Must be true for geometric connection with Peterson isomorphism.
- Corresponding result for  $s_\lambda^{(k)}$  is known, but this gives a Catalan/ $K$ -theoretic Catalan proof.
- $k$ -Rectangle Property fails for  $g_\lambda^{(k)}$ .

Summary:

Summary:

- $g_{\lambda}^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$  used for branching positivity.

Summary:

- $g_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$  used for branching positivity.
- $\tilde{g}_\lambda^{(k)} = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$  conjecturally related to  $K$ -Peterson isomorphism with many positivity conjectures.

Summary:

- $g_{\lambda}^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$  used for branching positivity.
- $\tilde{g}_{\lambda}^{(k)} = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$  conjecturally related to  $K$ -Peterson isomorphism with many positivity conjectures.

What can be said about  $K$ -theoretic Catalan functions in general?



# Positivity of $K$ -theoretic Catalan functions

Recall (Blasiak et al., 2020)

For  $\Psi$  any root ideal and  $\lambda$  a partition,  $H(\Psi; \lambda)$  is Schur positive.

# Positivity of $K$ -theoretic Catalan functions

Recall (Blasiak et al., 2020)

For  $\Psi$  any root ideal and  $\lambda$  a partition,  $H(\Psi; \lambda)$  is Schur positive.

Conjecture (Blasiak-Morse-S., 2020)

For  $\Psi$  a root ideal and  $\lambda$  a partition,

# Positivity of $K$ -theoretic Catalan functions

Recall (Blasiak et al., 2020)

For  $\Psi$  any root ideal and  $\lambda$  a partition,  $H(\Psi; \lambda)$  is Schur positive.

Conjecture (Blasiak-Morse-S., 2020)

For  $\Psi$  a root ideal and  $\lambda$  a partition,

- $K(\Psi; \Psi; \lambda) = \sum_{\mu} a_{\mu} g_{\mu}$  satisfies  $(-1)^{|\lambda|-|\mu|} a_{\mu} \in \mathbb{Z}_{\geq 0}$ .

# Positivity of $K$ -theoretic Catalan functions

## Recall (Blasiak et al., 2020)

For  $\Psi$  any root ideal and  $\lambda$  a partition,  $H(\Psi; \lambda)$  is Schur positive.

## Conjecture (Blasiak-Morse-S., 2020)

For  $\Psi$  a root ideal and  $\lambda$  a partition,

- $K(\Psi; \Psi; \lambda) = \sum_{\mu} a_{\mu} g_{\mu}$  satisfies  $(-1)^{|\lambda|-|\mu|} a_{\mu} \in \mathbb{Z}_{\geq 0}$ .
- $K(\Psi; RC^a(\Psi); \lambda) = \sum_{\mu} b_{\mu} s_{\mu}$  satisfies  $b_{\mu} \in \mathbb{Z}_{\geq 0}$ .

For  $G_\lambda^{(k)}$  an affine Grothendieck polynomial (dual to  $g_\lambda^{(k)}$ ),

For  $G_\lambda^{(k)}$  an affine Grothendieck polynomial (dual to  $g_\lambda^{(k)}$ ),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_{\mu} ?? g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_{\lambda} ?? G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

For  $G_\lambda^{(k)}$  an affine Grothendieck polynomial (dual to  $g_\lambda^{(k)}$ ),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_{\mu} ?? g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_{\lambda} ?? G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

- 2 Combinatorially describe branching coefficients:  $g_\lambda^{(k)} = \sum_{\mu} ?? g_\mu^{(k+1)}$ .

For  $G_\lambda^{(k)}$  an affine Grothendieck polynomial (dual to  $g_\lambda^{(k)}$ ),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_{\mu} ?? g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_{\lambda} ?? G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

- 2 Combinatorially describe branching coefficients:  $g_\lambda^{(k)} = \sum_{\mu} ?? g_\mu^{(k+1)}$ .

- 3 Combinatorially describe  $g_\lambda^{(k)} = \sum_{\mu} ?? s_\mu^{(k)}$ .



Raising operator techniques extend beyond Schubert calculus! Work by Blasiak-Haiman-Morse-Pun-S.:

- Shuffle theorems (Blasiak et al., 2021a; 2021b).
- Macdonald polynomials and LLT polynomials (Blasiak et al., 2021c).
- Much more work to be done!

## Thank you!

- Anderson, David, Linda Chen, and Hsian-Hua Tseng. 2017. *On the quantum  $K$ -ring of the flag manifold*, preprint. arXiv: 1711.08414.
- Blasiak, Jonah, Mark Haiman, Jennifer Morse, Anna Pun, and George H Seelinger. 2021a. *A Shuffle Theorem for Paths Under Any Line*, arXiv e-prints, available at arXiv:2102.07931.
- Blasiak, Jonah, Mark Haiman, Jennifer Morse, Anna Pun, and George H. Seelinger. 2021b. *A proof of the Extended Delta Conjecture*, arXiv e-prints, available at arXiv:2102.08815.
- . 2021c. *LLT polynomials in the Schiffmann algebra*. In preparation.
- Blasiak, Jonah, Jennifer Morse, Anna Pun, and Daniel Summers. 2019. *Catalan Functions and  $k$ -Schur Positivity*, J. Amer. Math. Soc. **32**, no. 4, 921–963.
- Blasiak, Jonah, Jennifer Morse, and Anna Pun. 2020. *Demazure crystals and the Schur positivity of Catalan functions*, preprint. arXiv: 2007.04952.
- Blasiak, Jonah, Jennifer Morse, and George H. Seelinger. 2020.  *$K$ -theoretic Catalan functions*, preprint. arXiv: 2010.01759.
- Chen, Li-Chung. 2010. *Skew-linked partitions and a representation theoretic model for  $k$ -Schur functions*, Ph.D. thesis.
- Fomin, Sergey, Sergei Gelfand, and Alexander Postnikov. 1997. *Quantum Schubert polynomials*, J. Amer. Math. Soc. **10**, no. 3, 565–596, DOI 10.1090/S0894-0347-97-00237-3. MR1431829
- Ikeda, Takeshi, Shinsuke Iwao, and Toshiaki Maeno. 2018. *Peterson Isomorphism in  $K$ -theory and Relativistic Toda Lattice*, preprint. arXiv: 1703.08664.
- Lam, Thomas. 2008. *Schubert polynomials for the affine Grassmannian*, J. Amer. Math. Soc. **21**, no. 1, 259–281.
- Lam, Thomas, Luc Lapointe, Jennifer Morse, and Mark Shimozono. 2010. *Affine insertion and Pieri rules for the affine Grassmannian*, Mem. Amer. Math. Soc. **208**, no. 977.
- Lam, Thomas, Anne Schilling, and Mark Shimozono. 2010.  *$K$ -theory Schubert calculus of the affine Grassmannian*, Compositio Math. **146**, 811–852.
- Lapointe, Luc, Alain Lascoux, and Jennifer Morse. 2003. *Tableau atoms and a new Macdonald positivity conjecture*, Duke Mathematical Journal **116**, no. 1, 103–146.
- Morse, Jennifer. 2011. *Combinatorics of the  $K$ -theory of affine Grassmannians*, Advances in Mathematics.
- Panyushev, Dmitri I. 2010. *Generalised Kostka-Foulkes polynomials and cohomology of line bundles on homogeneous vector bundles*, Selecta Math. (N.S.) **16**, no. 2, 315–342.

$$k_m^{(r)} = \sum_{i=0}^m \binom{r+i-1}{i} h_{m-i} = s_m(X+r),$$

a specialization of “multiSchur functions.” See, e.g., Lascoux-Naruse (2014).

$$k_\gamma = k_{\gamma_1}^{(0)} k_{\gamma_2}^{(1)} \dots k_{\gamma_\ell}^{(\ell-1)}$$